

Response function of strongly interacting Fermi gas in a virial expansion

Gang Shen*

*Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA
and Institute for Nuclear Theory, University of Washington, Seattle, WA 98195, USA*

(Dated: August 14, 2012)

The dynamic response functions of strongly interacting fermion gas in homogeneous space are investigated in a virial expansion to second order. The density response function exhibits transition from atomic to molecular response, as the interaction strength increases and the system undergoes BCS-BEC crossover. The qualitative features of density and spin response agree with measurements from the Bragg spectroscopy experiments. The virial response is exact at low density and high temperature, therefore providing a benchmark for many-body response.

PACS numbers: 03.75.Hh, 03.75.Ss, 05.30.Fk

I. INTRODUCTION

Recent years the advances in cold atom experiments have improved our understanding of thermodynamic and dynamic properties of strongly interacting systems. Particularly, the dynamic response function gives the response of the system under an excitation probe, therefore describing the inelastic scattering processes. Quite remarkably, the Bragg spectroscopy has been used to measure the dynamic and static density response functions for a strongly interacting cold Fermion system [1]. A smooth transition from atomic to molecular spectra, or from BCS to BEC regimes, is observed with a clear signature of pairing at and above unitarity, which provides a direct link to two-body correlations. Recently, the Bragg spectroscopy is also applied to obtain the dynamic spin response function [2].

Due to its nonperturbative nature, theoretically it is extremely hard to obtain the response function of strongly interacting Fermion system. There have been several investigations on the dynamic response function [3–7]. In certain limits, large momentum transfer between external probe and the system [8], or system being at very low density, to name a few, the many-body phenomena are expected to be dominated by few-body physics explicitly. The virial expansion presents a tractable approach to the strongly interacting system and has a controllable small parameter, the fugacity $z = \exp(\mu/T)$, when the system is at low density (chemical potential μ) and high temperature (T). It has been applied to study the thermodynamic properties of strongly interacting Fermi gas [9, 10]. For a trapped strongly interacting Fermi gas, a quantum virial expansion to second order for the response functions has been developed in Refs. [11, 12].

In this work, the latter virial expansion is extended to study the dynamic response functions of strongly interacting Fermi gas in homogeneous space. The formalism closely follows that developed in Ref. [11] for trapped Fermi gas. The dynamic density response function in homogeneous space is found to exhibit transition from atomic to molecular spectra, as the interaction strength increases and the system undergoes BCS-BEC crossover. The virial response is exact at low density and high temperature, when fugacity is a small number. Therefore it provides a benchmark for the many-body response functions. Qualitatively the response functions of strongly interacting fermi gas in homegeneous space show similar characteristics as those in trapped fermi gas. In fact they may be related by a local density approximation. In this work, the results in homogeneous space are explicitly written down in compact and closed form with simple integrals. The virial expansion for dynamic response function can be readily applied to other strongly interacting many-body system, like neutron matter and nuclear matter in supernova (see Ref. [13] for a study on the long wavelength behavior of static response function).

The paper is organized as follows. In next section II the formalism for the dynamic response function of Fermi gas in second order virial expansion is presented in detail. In Sec. III the frequency dependence, interaction dependence, and temperature dependence of dynamic response function, as well as the static response function are discussed. The conclusions are given in Sec. IV.

The natural units $\hbar = k_B = 1$ are used throughout.

*Electronic address: gshen@uw.edu

II. THE DYNAMIC DENSITY RESPONSE

The formalism below closely follows that developed in Ref. [11] for a trapped Fermi gas. For completeness the formalism is given in detail here. The dynamic density response for a spin-unpolarized gas under four momentum transfer ($\mathbf{q}, \omega$) can be separated into two pieces:

$$S_D(\mathbf{q}, \omega) = 2[S_{\uparrow\uparrow}(\mathbf{q}, \omega) + S_{\uparrow\downarrow}(\mathbf{q}, \omega)], \quad (2.1)$$

where $S_{\downarrow\downarrow} = S_{\uparrow\uparrow}$ and $S_{\downarrow\uparrow} = S_{\uparrow\downarrow}$ are used. Similarly one could obtain the dynamic spin response function,

$$S_S(\mathbf{q}, \omega) = 2[S_{\uparrow\uparrow}(\mathbf{q}, \omega) - S_{\uparrow\downarrow}(\mathbf{q}, \omega)]. \quad (2.2)$$

It is most convenient to relate the response function to a time-dependent correlation function in imaginary time, via analytic continuation and the fluctuation-dissipation theorem, *i.e.* the dynamic susceptibility,

$$\chi_{\sigma\sigma'}(\mathbf{r}, \mathbf{r}', \tau) = -\langle T_\tau \hat{n}_\sigma(\mathbf{r}, \tau) \hat{n}_{\sigma'}(\mathbf{r}, 0) \rangle \quad (2.3)$$

where T_τ is the imaginary-time ordering operator, and $\hat{n}_\sigma(\mathbf{r}, \tau)$ is the density (fluctuation) operator in spin channel σ , and τ is an imaginary time in the interval $0 < \tau \leq \beta = 1/k_B T$. With these definitions, one can show that

$$S_{\sigma\sigma'}(\mathbf{r}, \mathbf{r}', \omega) = \frac{-\text{Im}\chi_{\sigma\sigma'}(\mathbf{r}, \mathbf{r}'; i\omega_n \rightarrow \omega + i0^+)}{\pi(1 - e^{-\beta\omega})}, \quad (2.4)$$

where the matsubara frequencies $\omega_n = 2\pi n k_B T$ ($n = 0, \pm 1, \dots$). Above equation can be Fourier-transformed with respect to $\mathbf{r} - \mathbf{r}'$ to obtain $S_{\sigma\sigma'}(\mathbf{q}, \omega)$. By the usual virial expansion in fugacity $z = e^{\beta\mu}$,

$$\chi_{\sigma\sigma'} = zX_1[1 + z(X_2/X_1 - Q_1) + \dots], \quad (2.5)$$

where

$$X_N = -\text{Tr}_N \left[e^{-\beta\hat{H}} e^{\tau\hat{H}} \hat{n}_\sigma(\mathbf{r}) e^{-\tau\hat{H}} \hat{n}_{\sigma'}(\mathbf{r}') \right]. \quad (2.6)$$

The interaction enters from second order term. Using completeness relations,

$$\Delta\chi_{\sigma\sigma',2}(\mathbf{r}, \mathbf{r}', \tau) = \Delta X_2 = X_2 - X_2^0 = \quad (2.7)$$

$$= -\sum_{P,Q} \left[e^{-\beta E_P + \tau(E_P - E_Q)} C_{\sigma\sigma'}^{PQ}(\mathbf{r}, \mathbf{r}') \right], \quad (2.8)$$

where

$$C_{\sigma\sigma'}^{PQ}(\mathbf{r}, \mathbf{r}') = \langle P | \hat{n}_\sigma(\mathbf{r}) | Q \rangle \langle Q | \hat{n}_{\sigma'}(\mathbf{r}') | P \rangle \quad (2.9)$$

and pair state ($\uparrow\downarrow$) $|P\rangle$ ($|Q\rangle$) has energy E_P (E_Q). One can perform discrete Fourier transform on $\Delta\chi_{\sigma\sigma',2}(\mathbf{r}, \mathbf{r}', \tau)$, and analytical continue the results via $i\omega_n \rightarrow \omega + i\eta$, to get the second order response function,

$$\begin{aligned} \Delta S_{\sigma\sigma',2}(\mathbf{r}, \mathbf{r}', \omega) &= -\frac{\text{Im}\Delta\chi_{\sigma\sigma',2}(\mathbf{r}, \mathbf{r}', \omega)}{(1 - e^{-\beta\omega})\pi} \\ &= \sum_{P,Q} \delta(\omega + E_P - E_Q) e^{-\beta E_P} C_{\sigma\sigma'}^{PQ}(\mathbf{r}, \mathbf{r}'). \end{aligned} \quad (2.10)$$

The remaining step to get momentum space response function is through Fourier transform as follows,

$$\Delta S_{\sigma\sigma',2}(\mathbf{q}, \omega) = \int d\mathbf{R} e^{-i\mathbf{Q}\cdot\mathbf{R}} \int d\mathbf{r} d\mathbf{r}' e^{i\mathbf{r}(\mathbf{Q}_c/2 - \mathbf{q}) + i\mathbf{r}'(\mathbf{Q}_c/2 + \mathbf{q})} \Delta S_{\sigma\sigma',2}(\mathbf{r}, \mathbf{r}', \omega) \quad (2.11)$$

$$= \sum_{PQ} \delta(\omega + E_P - E_Q) e^{-\beta E_P} F_{\sigma\sigma'}^{PQ}(\mathbf{q}), \quad (2.12)$$

where

$$F_{\sigma\sigma'}^{PQ}(\mathbf{q}) = \int d\mathbf{r}d\mathbf{r}' e^{-i\mathbf{q}(\mathbf{r}-\mathbf{r}')} C_{\sigma\sigma'}^{PQ}(\mathbf{r}, \mathbf{r}'). \quad (2.13)$$

A reminder here for the form of $C_{\sigma\sigma'}^{PQ}(\mathbf{r}, \mathbf{r}')$,

$$\begin{aligned} C_{\uparrow\uparrow}^{PQ}(\mathbf{r}, \mathbf{r}') &= \int d\mathbf{r}_2 d\mathbf{r}_2' \Phi_P^*(\mathbf{r}, \mathbf{r}_2) \Phi_Q(\mathbf{r}, \mathbf{r}_2) \Phi_Q^*(\mathbf{r}', \mathbf{r}_2') \Phi_P(\mathbf{r}', \mathbf{r}_2'), \\ C_{\uparrow\downarrow}^{PQ}(\mathbf{r}, \mathbf{r}') &= \int d\mathbf{r}_1 d\mathbf{r}_2 \Phi_P^*(\mathbf{r}, \mathbf{r}_2) \Phi_Q(\mathbf{r}, \mathbf{r}_2) \Phi_Q^*(\mathbf{r}_1, \mathbf{r}') \Phi_P(\mathbf{r}_1, \mathbf{r}'), \end{aligned} \quad (2.14)$$

One can further decompose the response function in center of mass (C.M.) and relative parts, by making use of

$$\delta(\omega + E_P - E_Q) e^{-\beta E_P} = \int d\omega' \delta(\omega' + \epsilon_{p1} - \epsilon_{q1}) \delta(\omega - \omega' + \epsilon_{p2} - \epsilon_{q2}) e^{-\beta(\epsilon_{p1} + \epsilon_{p2})}, \quad (2.15)$$

where $\epsilon_{p1(q1)}, \epsilon_{p2(q2)}$ are the C.M. and relative part of energy.

Now one has to differentiate between $\uparrow\uparrow$ and $\uparrow\downarrow$ cases. For $\uparrow\uparrow$, one needs to group $(\mathbf{r}, \mathbf{r}_2)$, and $(\mathbf{r}', \mathbf{r}_2')$ separately. Specifically, define C.M. and relative parts as follows,

$$\begin{aligned} \mathbf{r} &= \mathbf{R} + \mathbf{x}/2, \\ \mathbf{r}_2 &= \mathbf{R} - \mathbf{x}/2, \end{aligned} \quad (2.16)$$

and do the same for $(\mathbf{r}', \mathbf{r}_2')$. For $\uparrow\downarrow$, one needs to group $(\mathbf{r}, \mathbf{r}_2)$, and $(\mathbf{r}_1, \mathbf{r}')$ separately. After some algebra, one can arrive at following forms,

$$\Delta S_{\sigma\sigma',2} = \left\{ \int d\omega' W_{cm}(\mathbf{q}, \omega') W_{rel}^{\sigma\sigma'}(\mathbf{q}, \omega - \omega') \right\}, \quad (2.17)$$

with following notations,

$$W_{cm} \equiv \sum_{p1q1} \delta(\omega' + \epsilon_{p1} - \epsilon_{q1}) e^{-\beta\epsilon_{p1}} |f_{p1q1}|^2, \quad (2.18)$$

where,

$$f_{p1q1} \equiv \int d\mathbf{R} e^{-i\mathbf{q}\cdot\mathbf{R}} \varphi_{p1}^*(\mathbf{R}) \varphi_{q1}(\mathbf{R}), \quad (2.19)$$

and

$$W_{rel}^{\sigma\sigma'} \equiv \sum_{p2q2} \delta(\omega - \omega' + \epsilon_{p2} - \epsilon_{q2}) e^{-\beta\epsilon_{p2}} A_{p2q2}^{\sigma\sigma'}, \quad (2.20)$$

with $A_{p2q2}^{\uparrow\uparrow} \equiv |A_{p2q2}|^2$ and $A_{p2q2}^{\uparrow\downarrow} \equiv A_{p2q2} A_{q2p2}$, and

$$A_{p2q2} \equiv \int d\mathbf{x} e^{-i\mathbf{q}\cdot\mathbf{x}/2} \phi_{p2}^*(\mathbf{x}) \phi_{q2}(\mathbf{x}). \quad (2.21)$$

The center of mass piece is trivial,

$$W_{cm}(\omega') \equiv \frac{m^2}{n\pi^2\beta q} e^{-(\omega' - q^2/4m)^2 \beta m/q^2}, \quad (2.22)$$

where the number of particles has been divided. Next the relative piece is discussed in detail, from BCS to BEC side.

A. BCS side

Using the scattering state wave function $\phi_S(px)$ in Eq. (A1), the relative piece in Eq. (2.20) can be written as,

$$W_{rel}^{\sigma\sigma'}(q, \omega) = \int dp_2 p_2^2 dq_2 q_2^2 \delta(\omega + \epsilon_{p_2} - \epsilon_{q_2}) e^{-\beta \epsilon_{p_2}} \left[\left(\int dx x^2 j_0(qx/2) \phi_S(p_2x) \phi_S(q_2x) \right)^2 \right. \\ \left. + \sum_{l>0} (2l+1)(-1)^{l(1-\delta_{\sigma\sigma'})} \left(\int dx x^2 j_l(qx/2) \sqrt{\frac{2}{\pi}} j_l(p_2x) \phi_S(q_2x) \right)^2 \right. \\ \left. + \sum_{l>0} (2l+1)(-1)^{l(1-\delta_{\sigma\sigma'})} \left(\int dx x^2 j_l(qx/2) \sqrt{\frac{2}{\pi}} j_l(q_2x) \phi_S(p_2x) \right)^2 - \text{non. inter. terms} \right]. \quad (2.23)$$

Note either $\phi_S(p_2x)$ or $\phi_S(q_2x)$ has to be interacting S wave state, otherwise it will cancel with corresponding non-interacting term. With above equation and Eq. (2.22) one can proceed to the dynamic structure function in Eq. (2.17),

$$\Delta \tilde{S}_{\sigma\sigma',2}(q, \omega) = \int d\tilde{p}_2 \tilde{p}_2^2 d\tilde{q}_2 \tilde{q}_2^2 \frac{3\tilde{T}}{4\tilde{q}} e^{-(\tilde{\omega} + 2\tilde{p}_2^2 - 2\tilde{q}_2^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} - 2\tilde{p}_2^2 / \tilde{T}} \left[\left(\int dx x^2 j_0(\tilde{q}x/2) \phi_S(\tilde{p}_2x) \phi_S(\tilde{q}_2x) \right)^2 \right. \\ \left. + \sum_{l>0} (2l+1)(-1)^{l(1-\delta_{\sigma\sigma'})} \left(\int dx x^2 j_l(\tilde{q}x/2) \sqrt{\frac{2}{\pi}} j_l(\tilde{p}_2x) \phi_S(\tilde{q}_2x) \right)^2 \right. \\ \left. + \sum_{l>0} (2l+1)(-1)^{l(1-\delta_{\sigma\sigma'})} \left(\int dx x^2 j_l(\tilde{q}x/2) \sqrt{\frac{2}{\pi}} j_l(\tilde{q}_2x) \phi_S(\tilde{p}_2x) \right)^2 - \text{non. inter. terms} \right], \quad (2.24)$$

where scaled variables, momentum in terms of k_F and energy in terms of $\epsilon_F = k_F^2/2m$, have been used. For instance, $\tilde{q} = q/k_F$, $\tilde{T} = T/\epsilon_F$, and $\Delta \tilde{S}_{\sigma\sigma',2} = \Delta S_{\sigma\sigma',2} \times \epsilon_F$.

a) The first term, S wave to S wave scattering, (and its non-interacting counterpart) in Eq. (2.24) can be worked out as follows,

$$\Delta \tilde{S}_{\sigma\sigma',2}^a(q, \omega) = \int d\tilde{p}_2 d\tilde{q}_2 \frac{3\tilde{T}}{4\tilde{q}^3} \left(\frac{2}{\pi} \right)^2 e^{-(\tilde{\omega} + 2\tilde{p}_2^2 - 2\tilde{q}_2^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} - 2\tilde{p}_2^2 / \tilde{T}} \times \\ \left[\frac{\pi}{4} C_{p2} C_{q2} (-\text{sign}(\tilde{q}/2 - \tilde{p}_2 - \tilde{q}_2) + \text{sign}(\tilde{q}/2 + \tilde{p}_2 - \tilde{q}_2) + \text{sign}(\tilde{q}/2 - \tilde{p}_2 + \tilde{q}_2) - \text{sign}(\tilde{q}/2 + \tilde{p}_2 + \tilde{q}_2)) \right. \\ \left. + \frac{\pi}{4} S_{p2} S_{q2} (\text{sign}(\tilde{q}/2 - \tilde{p}_2 - \tilde{q}_2) + \text{sign}(\tilde{q}/2 + \tilde{p}_2 - \tilde{q}_2) + \text{sign}(\tilde{q}/2 - \tilde{p}_2 + \tilde{q}_2) + \text{sign}(\tilde{q}/2 + \tilde{p}_2 + \tilde{q}_2)) \right. \\ \left. + \frac{1}{4} C_{p2} S_{q2} \log \frac{(1+v_2)^2}{(1-v_2)^2} + \frac{1}{4} C_{q2} S_{p2} \log \frac{(1+v_3)^2}{(1-v_3)^2} \right]^2 - \text{non. inter. terms}, \quad (2.25)$$

where $\text{sign}()$ is the sign function, and following quantities are defined,

$$C_{p2} = \frac{1}{\sqrt{1 + \tilde{p}_2^2 k_F^2 a^2}}, \quad (2.26)$$

$$S_{p2} = \frac{-\tilde{p}_2 k_F a}{\sqrt{1 + \tilde{p}_2^2 k_F^2 a^2}}, \quad (2.27)$$

$$v_2 = \frac{\tilde{q}^2/4 + \tilde{p}_2^2 - \tilde{q}_2^2}{\tilde{q}\tilde{p}_2}, \quad (2.28)$$

$$v_3 = \frac{\tilde{q}^2/4 + \tilde{q}_2^2 - \tilde{p}_2^2}{\tilde{q}\tilde{q}_2}. \quad (2.29)$$

b) The second and third terms in Eq. (2.24) due to transition between $l > 0$ partial waves and S wave are simplified as follows,

$$\Delta \tilde{S}_{\sigma\sigma',2}^b(q, \omega) = \int d\tilde{p}_2 d\tilde{q}_2 \frac{3\tilde{T}}{4\tilde{q}^3} \left(\frac{2}{\pi} \right)^2 e^{-(\tilde{\omega} + 2\tilde{p}_2^2 - 2\tilde{q}_2^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} - 2\tilde{p}_2^2 / \tilde{T}} \sum_{l>0} (2l+1)(-1)^{l(1-\delta_{\sigma\sigma'})} \times$$

$$\left[\left(\frac{\pi}{2} \right)^2 P_l(v_2)^2 (C_{q_2}^2 - 1) + Q_l(v_2)^2 S_{q_2}^2 + \pi P_l(v_2) Q_l(v_2) C_{q_2} S_{q_2} \right. \\ \left. + \left(\frac{\pi}{2} \right)^2 P_l(v_3)^2 (C_{p_2}^2 - 1) + Q_l(v_3)^2 S_{p_2}^2 + \pi P_l(v_3) Q_l(v_3) C_{p_2} S_{p_2} \right], \quad (2.30)$$

where P_l and Q_l are l^{th} -order Legendre polynomials of first and second kind, respectively. Note except $(Q_l)^2$ terms, the other terms are nonzero only for $|v_2|$ or $|v_3| \leq 1$ [14]. Using the summation relation for the product of two Legendre polynomials (see Appendix B), b) term can be further simplified as follows.

1. Spin anti-parallel case

$$\Delta \tilde{S}_{\uparrow\downarrow,2}^b(q, \omega) = \int d\tilde{p}_2 d\tilde{q}_2 \frac{3\tilde{T}}{4\tilde{q}^3} \left(\frac{2}{\pi} \right)^2 e^{-(\tilde{\omega} + 2\tilde{p}_2^2 - 2\tilde{q}_2^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} - 2\tilde{p}_2^2 / \tilde{T}} \times \\ \left[\left(\frac{\pi}{2} \right)^2 (\delta(v_2) - 1) (C_{q_2}^2 - 1) + \left(\frac{1}{4v_2} \log \left(\frac{1+v_2}{1-v_2} \right)^2 - Q_0(v_2)^2 \right) S_{q_2}^2 + \pi \left(\frac{1}{2v_2} - Q_0(v_2) \right) C_{q_2} S_{q_2} \right. \\ \left. + \left(\frac{\pi}{2} \right)^2 (\delta(v_3) - 1) (C_{p_2}^2 - 1) + \left(\frac{1}{4v_3} \log \left(\frac{1+v_3}{1-v_3} \right)^2 - Q_0(v_3)^2 \right) S_{p_2}^2 + \pi \left(\frac{1}{2v_3} - Q_0(v_3) \right) C_{p_2} S_{p_2} \right]. \quad (2.31)$$

Note again the 1st, 3rd, 4th, and 6th terms are nonzero only for $|v_2|$ or $|v_3| \leq 1$. Principal-value integrals are assumed where it is appropriate.

2. Spin parallel case

$$\Delta \tilde{S}_{\uparrow\uparrow,2}^b(q, \omega) = \int d\tilde{p}_2 d\tilde{q}_2 \frac{3\tilde{T}}{4\tilde{q}^3} \left(\frac{2}{\pi} \right)^2 e^{-(\tilde{\omega} + 2\tilde{p}_2^2 - 2\tilde{q}_2^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} - 2\tilde{p}_2^2 / \tilde{T}} \times \\ \left[- \left(\frac{\pi}{2} \right)^2 P_0(v_2) (C_{q_2}^2 - 1) + \left(\frac{1}{v_2^2 - 1} - Q_0(v_2)^2 \right) S_{q_2}^2 - \pi P_0(v_2) Q_0(v_2) C_{q_2} S_{q_2} \right. \\ \left. - \left(\frac{\pi}{2} \right)^2 P_0(v_3) (C_{p_2}^2 - 1) + \left(\frac{1}{v_3^2 - 1} - Q_0(v_3)^2 \right) S_{p_2}^2 - \pi P_0(v_3) Q_0(v_3) C_{p_2} S_{p_2} \right]. \quad (2.32)$$

Similarly, the 1st, 3rd, 4th, and 6th terms are nonzero only for $|v_2|$ or $|v_3| \leq 1$.

Summing up Eqs. (2.25) with (2.31) or (2.32), one could obtain the dynamic response functions on the BCS side as follows,

$$\tilde{S}_{\uparrow\downarrow}^{BCS}(q, \omega) = z^2 (\Delta \tilde{S}_{\uparrow\downarrow,2}^a(q, \omega) + \Delta \tilde{S}_{\uparrow\downarrow,2}^b(q, \omega)), \quad (2.33)$$

$$\tilde{S}_{\uparrow\uparrow}^{BCS}(q, \omega) = \tilde{S}_F(q, \omega) + z^2 (\Delta \tilde{S}_{\uparrow\uparrow,2}^a(q, \omega) + \Delta \tilde{S}_{\uparrow\uparrow,2}^b(q, \omega)), \quad (2.34)$$

where $\tilde{S}_F(q, \omega)$ is the dynamic response function for non-interacting Fermi gas [15],

$$\tilde{S}_F(q, \omega) = \frac{3\tilde{T}}{16\tilde{q}} \frac{1}{1 - e^{-\tilde{\omega}/\tilde{T}}} \log \left[\frac{1 + ze^{-(\tilde{\omega}/\tilde{q} - \tilde{q})^2/4\tilde{T}}}{1 + ze^{-(\tilde{\omega}/\tilde{q} + \tilde{q})^2/4\tilde{T}}} \right]. \quad (2.35)$$

Note a factor of half is included in above equation to remove spin degeneracy.

B. BEC side

On BEC side there exist contributions from bound state $\phi_b(x)$ in Eq. (A2), besides scattering states.

c) Transition between bound state and scattering state

$$\Delta \tilde{S}_{\sigma\sigma',2}^c(q, \omega) = \int d\tilde{q}_2 \tilde{q}_2^2 \frac{3\tilde{T}}{4\tilde{q}} e^{-(\tilde{\omega} - 2/k_F^2 a^2 - 2\tilde{q}_2^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} + 2/\tilde{T} k_F^2 a^2} \left[\left(\int dx x^2 j_0(\tilde{q}x/2) \phi_S(\tilde{q}_2 x) \sqrt{4\pi} \phi_b(x) \right)^2 \right. \\ \left. + \sum_{l>0} (2l+1) (-1)^{l(1-\delta_{\sigma\sigma'})} \left(\int dx x^2 j_l(\tilde{q}x/2) \sqrt{\frac{2}{\pi}} j_l(\tilde{q}_2 x) \sqrt{4\pi} \phi_b(x) \right)^2 \right] \\ + \int d\tilde{p}_2 \tilde{p}_2^2 \frac{3\tilde{T}}{4\tilde{q}} e^{-(\tilde{\omega} + 2\tilde{p}_2^2 + 2/k_F^2 a^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} - 2\tilde{p}_2^2 / \tilde{T}} \left[\left(\int dx x^2 j_l(\tilde{q}x/2) \phi_S(\tilde{p}_2 x) \sqrt{4\pi} \phi_b(x) \right)^2 \right]$$

$$\begin{aligned}
& + \sum_{l>0} (2l+1)(-1)^{l(1-\delta_{\sigma\sigma'})} \left(\int dx x^2 j_l(\tilde{q}x/2) \sqrt{\frac{2}{\pi}} j_l(\tilde{p}_2 x) \sqrt{4\pi} \phi_b(x) \right)^2 \Big] \\
& = \int d\tilde{q}_2 \frac{3\tilde{T}}{4\tilde{q}^3} e^{-(\tilde{\omega}-2/k_F^2 a^2 - 2\tilde{q}_2^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} + 2/\tilde{T} k_F^2 a^2} \left[(A(u_2) - Q_0(u_2)^2) \right. \\
& \quad + (Q_0(u_2)^2 C_{q_2} + (\arctan[(\tilde{q}/2 - \tilde{q}_2)k_F a] + \arctan[(\tilde{q}/2 + \tilde{q}_2)k_F a]) S_{q_2})^2 \\
& \quad + \int d\tilde{p}_2 \frac{3\tilde{T}}{4\tilde{q}^3} e^{-(\tilde{\omega}+2\tilde{p}_2^2+2/k_F^2 a^2 - \tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} - 2\tilde{p}_2^2/\tilde{T}} \left[(A(u_3) - Q_0(u_3)^2) \right. \\
& \quad \left. + (Q_0(u_3)^2 C_{p_2} + (\arctan[(\tilde{q}/2 - \tilde{p}_2)k_F a] + \arctan[(\tilde{q}/2 + \tilde{p}_2)k_F a]) S_{p_2})^2 \right], \tag{2.36}
\end{aligned}$$

where the following notations have been used,

$$u_2 = \frac{1/k_F^2 a^2 + \tilde{q}^2/4 + \tilde{q}_2^2}{\tilde{q}\tilde{q}_2}, \tag{2.37}$$

$$u_3 = \frac{1/k_F^2 a^2 + \tilde{q}^2/4 + \tilde{p}_2^2}{\tilde{q}\tilde{p}_2}, \tag{2.38}$$

$$A(x) = \frac{1}{x^2 - 1}, \quad \text{for spin } \uparrow\uparrow, \tag{2.39}$$

$$A(x) = Q_0(1/x)/x, \quad \text{for spin } \uparrow\downarrow. \tag{2.40}$$

d) Transition between bound states, or molecular response

$$\begin{aligned}
\Delta\tilde{S}_{\sigma\sigma',2}^d(q,\omega) &= \frac{3\tilde{T}}{4\tilde{q}} e^{-(\tilde{\omega}-\tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} + 2/\tilde{T} k_F^2 a^2} \left[\int dx x^2 j_0(\tilde{q}x/2) 4\pi \phi_b(x)^2 \right]^2 \\
&= \frac{3\pi\tilde{T}}{2\tilde{q}^2 k_F a} e^{-(\tilde{\omega}-\tilde{q}^2/2)^2 / 2\tilde{q}^2 \tilde{T} + 2/\tilde{T} k_F^2 a^2} (1 + \tilde{q}^2 k_F^2 a^2)^{-1/2} P_{-1/2}^{-1/2} [(1 + \tilde{q}^2 k_F^2 a^2)^{-1/2}]^2. \tag{2.41}
\end{aligned}$$

It is clear from Eq. (2.41) that on deep BEC side, $k_F a \ll 1$, the molecular response dominates in the dynamic response function.

Summing up Eqs. (2.25) with (2.31) or (2.32), as well as (2.36) and (2.41), one could obtain the dynamic response functions on the BEC side as follows,

$$\tilde{S}_{\uparrow\downarrow}^{BEC}(q,\omega) = z^2 (\Delta\tilde{S}_{\uparrow\downarrow,2}^a(q,\omega) + \Delta\tilde{S}_{\uparrow\downarrow,2}^b(q,\omega) + \Delta\tilde{S}_{\uparrow\downarrow,2}^c(q,\omega) + \Delta\tilde{S}_{\uparrow\downarrow,2}^d(q,\omega)), \tag{2.42}$$

$$\tilde{S}_{\uparrow\uparrow}^{BEC}(q,\omega) = \tilde{S}_F(q,\omega) + z^2 (\Delta\tilde{S}_{\uparrow\uparrow,2}^a(q,\omega) + \Delta\tilde{S}_{\uparrow\uparrow,2}^b(q,\omega) + \Delta\tilde{S}_{\uparrow\uparrow,2}^c(q,\omega) + \Delta\tilde{S}_{\uparrow\uparrow,2}^d(q,\omega)), \tag{2.43}$$

III. RESULTS AND DISCUSSIONS

In this section, the dependences of dynamic response function on frequency, interaction, and temperature, as well as static response function are discussed in details. The Bragg spectroscopy experiments have been carried out using large momentum transfer. Therefore in this section, the response functions are calculated with a large momentum transfer $q = 3k_F$.

A. Frequency dependence of $S_{\uparrow\uparrow}$ and $S_{\uparrow\downarrow}$

In Figure 1, the dynamic response functions for spin-parallel and spin-anti-parallel cases, $S_{\uparrow\uparrow}$ and $S_{\uparrow\downarrow}$, are shown as function of frequency at different interaction strengths for $T = 2T_F$ (left panels) and $T = 3T_F$ (right panels). In all the results, the largest fugacity is $z=0.25$ (BCS side $1/k_F a = -1$ and $T/T_F=2$). These small values suggest that higher order corrections to the virial expansion will be small. $\omega_R = q^2/2m$ is the atomic recoil frequency where m is the atomic mass.

The effect of interaction is most clear in the dynamic response function for spin-anti-parallel case, $S_{\uparrow\downarrow}$, as Eqs. (2.33, 2.42) contain only interaction induced contributions. On the BCS side $1/k_F a = -1$, $S_{\uparrow\downarrow}$ is peaked around the molecular recoil frequency $\omega_R/2$ due to strong pair correlation from attractive interaction. The peak becomes more visible as temperature decreases and/or interactions strength increases. On the BEC side, the response is sharply

peaked around the molecular recoil frequency due to the formation of bound pair state, which dominates the response function. The response function $S_{\uparrow\uparrow}$, given in Eqs. (2.34, 2.43), receives contribution from non-interaction terms Eq. (2.35), which peaks at the atomic recoil frequency. On the BCS side, the interaction is too weak so the response is dominated by the non-interacting terms. On the BEC side, the peak of response function $S_{\uparrow\uparrow}$ moves toward the molecular recoil frequency as $S_{\uparrow\downarrow}$.




FIG. 1: (Color online) $S_{\uparrow\uparrow}(q=3k_F, \omega)$ and $S_{\uparrow\downarrow}(q=3k_F, \omega)$ as function of frequency at different $1/k_F a$ and at $T = 2T_F$ (left) and $T = 3T_F$ (right).

B. Interaction dependence of dynamic response function

In Figure 2 the dynamic density (left) and spin (right) responses at $T = 3T_F$, obtained by Eqs. (2.1, 2.2), are shown for various interaction strengths. The density response function peaks at atomic recoil frequency on BCS side. As interaction increases, the peak becomes red-shifted. On BEC side, the density response becomes peaked around molecular frequency. These features qualitatively agree with recent Bragg spectroscopy experiment [2], although the latter experiment was performed at extremely low temperature in the superfluid phase. The spin response has broad peak around atomic frequency both on BEC and BCS sides, which again agrees with the experiment [2].




FIG. 2: (Color online) Interaction dependence of dynamic density (left) and spin (right) responses at $T = 3T_F$.

C. Temperature dependence of dynamic response function

In Figure 3, the dynamic density and spin responses at unitarity are shown at different temperatures. The density response has a relatively narrow peak around $\omega_R/2$ at $T = 2T_F$, and becomes broadened and shifted to higher energy as temperature increases when non-interacting term becomes increasingly more important. On the other hand, the spin response peaks around ω_R .

D. Static response function

The static response functions are obtained via following integrals,

$$S_{\sigma\sigma'}(q) = 2 \int_{-\infty}^{\infty} d\omega S_{\sigma\sigma'}(q, \omega). \quad (3.1)$$

In Figure 4 the static response functions $S_{\uparrow\uparrow}(q=3k_F)$ and $S_{\uparrow\downarrow}(q=3k_F)$ are shown as function of $1/k_F a$ at $T = 3T_F$. $S_{\uparrow\uparrow}(q=3k_F)$ increases from about one in deep BEC side when $1/k_F a$ reduces, and it decreases quickly approaching unitarity. The rise and fall is due to contribution from transition between bound state and scattering state on the BEC side in Eq. (2.36), which vanishes at unitarity. On the other hand, $S_{\uparrow\downarrow}(q=3k_F)$ changes more smoothly from BEC to BCS side. Note at unitarity when $T = 3T_F$, the fugacity is about 0.13, therefore one expects the higher order terms contribute at about 10-20% of second order term displayed here.




FIG. 3: (Color online) Temperature dependence of dynamic density and spin responses at unitarity.

IV. CONCLUSIONS

In this work, the dynamic response functions of strongly interacting Fermi gas in homogeneous space are investigated in a virial expansion to second order. The dynamic density response function is found to exhibit transition from atomic response to molecular response, as the interaction strength increases and the system undergoes BCS-BEC crossover. The spin response function has a broad peak around atomic recoil frequency at high temperature. These features qualitatively agree with recent experiments via Bragg spectroscopy. The virial response is exact at low density and high temperature, when fugacity is a small number. The fugacity discussed in this work is relatively small $z \leq 0.25$. Therefore it provides a benchmark for the many-body response functions.

Qualitatively, the response functions of strongly interacting fermi gas in homegeneous space show similar characteristics as those in trapped fermi gas. In fact they may be related by a local density approximation, which would be interesting to study in future work.

To second order virial expansion, the dynamic response function in homogeneous space can be explicitly written down in compact and closed form with simple integrals. This illustrates that the virial expansion to response function may be also applied to other strongly interacting many-body system at low density and high temperature, like neutron matter and nuclear matter in supernova.

It is a pleasure to acknowledge the encouragement from Joe Carlson, Chuck Horowitz, and Sanjay Reddy, as well as discussions with C.-C. Chien, J. Drut, S. Gandolfi, and Y. Nishida. The work is supported by a grant from the DOE under contract DE-AC52-06NA25396 and the DOE topical collaboration to study “Neutrinos and nucleosynthesis in hot and dense matter”.




FIG. 4: (Color online) Static response functions $S_{\uparrow\uparrow}(q=3k_F)$ and $S_{\uparrow\downarrow}(q=3k_F)$ as function of $1/k_F a$ at $T = 3T_F$.

Appendix A: S-wave pair states in a $\delta^3(\vec{r})$ potential

The S wave scattering state is known to be,

$$\phi_S(pr) = \sqrt{\frac{2}{\pi(1+p^2a^2)}} \frac{\sin(pr) - pa \cos(pr)}{pr} Y_{00} = \sqrt{\frac{2}{\pi}} \frac{\sin(pr + \eta_p)}{pr} Y_{00}, \quad (\text{A1})$$

when η_p equals zero, it reduces to non-interacting S wave state. To be precise, there should be a factor $-\text{sign}(a)$. However this overall phase does not influence the results in the paper. The bound state on the BEC side ($a > 0$) is,

$$\phi_b(\mathbf{r}) = \frac{1}{\sqrt{2\pi a}} \frac{e^{-r/a}}{r}, \quad (\text{A2})$$

with the binding energy,

$$E_b = -\frac{1}{ma^2}. \quad (\text{A3})$$

Appendix B: Sum of product of two Legendre Polynomials

The infinite sum over product of two Legendre polynomials can be carried out utilizing following relations,

$$\sum_{l \geq 0} (-1)^l (2l+1) P_l(x)^2 = \delta(x), \quad x \in [-1, 1], \quad (\text{B1})$$

$$\sum_{l \geq 0} (2l+1) P_l(x) Q_l(x) = 0, \quad (\text{B2})$$

$$\sum_{l \geq 0} (-1)^l (2l+1) P_l(x) Q_l(x) = \frac{1}{2x}, \quad x \in [-1, 1], \quad (\text{B3})$$

$$\sum_{l \geq 0} (2l+1) Q_l(x)^2 = \left(\frac{\pi}{2}\right)^2 \sum_{l \geq 0} (2l+1) P_l(x)^2 + \frac{1}{x^2 - 1}, \quad (\text{B4})$$

$$\sum_{l \geq 0} (-1)^l (2l+1) Q_l(x)^2 = \frac{1}{2x} \log \left| \frac{1+x}{1-x} \right|. \quad (\text{B5})$$

Note in Eq. (B4) the two sums are separately divergent when $x \in (-1, 1)$, but their difference is finite.

Appendix C: Virial expansion of density

On BCS side, the following equation for number density,

$$\frac{n\lambda^3}{2} = \frac{(4\pi/\tilde{T})^{3/2}}{6\pi^2} = \frac{2}{\sqrt{\pi}} \int_0^\infty dt \frac{\sqrt{t}}{1+e^t/z} + \sqrt{2} z^2 e^{2/\tilde{T}(k_F a)^2} \text{Erfc}\left[-\sqrt{\frac{2}{\tilde{T}}} \frac{1}{k_F a}\right], \quad (\text{C1})$$

is used to solve fugacity z .

On BEC side, the equation for number density is changed accordingly due to the additional contribution from bound state,

$$\frac{n\lambda^3}{2} = \frac{(4\pi/\tilde{T})^{3/2}}{6\pi^2} = \frac{2}{\sqrt{\pi}} \int_0^\infty dt \frac{\sqrt{t}}{1+e^t/z} + \sqrt{2} z^2 e^{2/\tilde{T}(k_F a)^2} \left(2 - \text{Erfc}\left[\sqrt{\frac{2}{\tilde{T}}} \frac{1}{k_F a}\right] \right). \quad (\text{C2})$$

-
- [1] G. Veeravalli, E. Kuhnle, P. Dyke, and C. J. Vale, Phys. Rev. Lett. **101**, 250403 (2008).
 - [2] S. Hoinka, M. Lingham, M. Delehay, and C. J. Vale, arXiv:1203.4657.
 - [3] A. Minguzzi, G. Ferrari, and Y. Castin, Eur. Phys. J. D **17**, 49 (2001).
 - [4] H. P. Büchler, P. Zoller, and W. Zwerger, Phys. Rev. Lett. **93**, 080401 (2004).
 - [5] G. M. Bruun, and G. Baym, Phys. Rev. A **74**, 033623 (2006).
 - [6] S. Stringari, Phys. Rev. Lett. **102**, 110406 (2009).
 - [7] H. Guo, C.-C. Chien, and K. Levin, Phys. Rev. Lett. **105**, 120401 (2010).
 - [8] Y. Nishida, arXiv:1110.5926v3.
 - [9] T.-L. Ho and E. J. Mueller, Phys. Rev. Lett. **92**, 160404 (2004).
 - [10] X.-J. Liu, H. Hu, and P. D. Drummond, Phys. Rev. Lett. **102**, 160401 (2009).
 - [11] H. Hu, X.-J. Liu, and P. D. Drummond, Phys. Rev. A **81**, 033630 (2010).
 - [12] H. Hu and X.-J. Liu, Phys. Rev. A **85**, 023612 (2012).
 - [13] C. J. Horowitz and A. Schwenk, Phys. Lett. B **642**, 326 (2006).
 - [14] To be precise, one has to multiply a factor of $\frac{1}{2}$ when $v_2(v_3) = 1$ in front of $P_l(v_2)$. It has negligible influence on the integral.
 - [15] F. Mazzanti and A. Polls, Phys. Lett. A **263**, 416 (1999).